

ALP Soil ID	State/Prov.	Cty	Crop	Soil Series	Soil Classification
SRS-2405	BC	Phantom Creek Vineyard	vineyard	Typic Xeropsamment	
SRS-2404	WA	Grant	Org Potatoes	Shano silt loam, 2 to 5 percent slopes	Fluvialglacial
SRS-2403	NC	Bertie	soybean	Lynchburg sandy loam, loamy fine sand	Coarse-silty, mixed, superactive, mesic Xeric Haplocambids
SRS-2402	PA	Centre Cty	corn	Hagerstown silt loam, 3 to 8 percent slopes	Fine-loamy, siliceous, semiactive, thermic Aeric Paleaquults
SRS-2401	SD	Minnehaha	corn	Nora-Crofton complex, 6 to 9 percent slopes	Fine, mixed, semiactive, mesic Typic Hapludalfs
SRS-2315	WY	Crook	Pasture	Nevee silt loam, 6 to 10 percent slopes, eroded	Fine-silty, mixed, superactive, mesic Udic Haplustolls
SRS-2314	BC	Dry Creek Vineyard	vineyard	Aridic Haploxeroll	Coarse-silty, mixed, superactive, calcareous, mesic Aridic Ustorthents
SRS-2313	NE	Holt	rye / potato	Pivot loamy sand 0 - 1 % slopes	Fluvial Fan
SRS-2312	CT	Litchfield Cty	corn	Winooski silt loam	Sandy, mixed, mesic Entic Haplustolls
SRS-2311	SC	Anderson	Soybean	Cecil clay loam, 6 to 10 percent slopes, eroded	Coarse-silty, mixed, superactive, mesic Fluvaquentic Dystrudepts
SRS-2310	MT	Cascade	fallow	Ipano-Ticell loams, 0 to 4 percent slopes	Fine, kaolinitic, thermic Typic Kanhapludults
SRS-2309	IA	Boone	grass	Hayden-Storden loams, 25 to 50 percent slopes	Fine-loamy, mixed, superactive, frigid Typic Calcicustolls
SRS-2308	IN	Cass	Soybeans	Gilford fine sandy loam	Fine-loamy, mixed, superactive, mesic Typic Natrargids
SRS-2307	VA	Chesterfiled	Pasture	Mayodan sandy loam, clayey substratum, 2 to 6 percent slopes	Coarse-loamy, mixed, superactive, mesic Typic Endoaquolls
SRS-2306	SK		Wheat	Mainly Orthic Me sandy loam	Fine, mixed, semiactive, thermic Typic Hapludults
SRS-2305	IA	Cass	Corn	Marshall silty clay loam	
SRS-2304	AR	Washington	fallow	Pickwick silt loam, 3 to 8 percent slopes, eroded	
SRS-2303	MD	Caroline	corn	Hambrook sandy loam, 0 to 2 percent slopes	Fine-silty, mixed, semiactive, thermic Typic Paleudults
SRS-2302	OR	Morrow	Corn	Quincy loamy fine sand, 2 to 12 percent slopes	Fine-loamy, siliceous, semiactive, mesic Typic Hapludults
SRS-2301	ON	Huron	wheat		
SRS-2215	CA	Sonoma	Grapes	Goldridge fine sandy loam, 9 to 15 percent slopes	
SRS-2214	MI	Clinton	Pasture	Capac-Marlette loams, 1 to 6 percent slopes	Fine-loamy, mixed, superactive, isomesic Typic Haplustults
SRS-2213	ID	Power	Potato	Declo loam, saline-alkali, 0 to 2 percent slopes	
SRS-2212	NC	Rowan	Pasture	Cecil sandy clay loam, 8 to 15 percent slopes, moderately eroded	Coarse-loamy, mixed, superactive, mesic Xeric Haplocalcids
SRS-2211	BC	Black SageRd	vineyard	Typic Xeropsamment	Fine, kaolinitic, thermic Typic Kanhapludults
SRS-2210	PA	Wayne	Veg	Volusia channery silt loam, 3 to 8 percent slopes	Fine-loamy, mixed, active, mesic Aeric Fragiaquepts
SRS-2209	NE	Madison	corn	Thurman loamy fine sand, 0-2% slopes	Sandy, mixed, mesic Udorthentic Haplustolls
SRS-2208	NM	Eddy	pasture, Tall wheatgrass	Harkey very fine sandy loam, 0 to 1 percent slopes	Coarse-silty, mixed, superactive, calcareous, thermic Typic Torrfluvents
SRS-2207	TN	Davidson	Forest	Maury-Urban land complex, 2 to 7 percent slopes	Fine, mixed, active, mesic Typic Paleudalfs
SRS-2206	AB		Canola		orthic dark brown chernozems
SRS-2205	CA	Sacramento	Grapes	Bruella sandy loam, 2 to 5 percent slopes	Fine-loamy, mixed, active, thermic Ultic Palexeralfs
SRS-2204	IA	Worth	corn	Webster clay loam, 0 to 2 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Endoaquolls
SRS-2203	MT	Yellowstone	wheat/fallow	Danvers-Shaak clay loams, 7 to 15 percent slopes	Fine, smectitic, frigid Vertic Argiustolls
SRS-2202	WV	Rallich	Pasture	Cotaco loam, 3 to 8 percent slopes	Fine-loamy, mixed, semiactive, mesic Aquic Hapludults
SRS-2201	PE	Queens	cover crop		
SRS-2115	OK	Payne	fallow	Norge loam, 1 to 3 percent slopes	Fine-silty, mixed, active, thermic Udic Paleustolls
SRS-2114	SK		Small Grains	Weyburn Clay Loam	Clay Loam
SRS-2113	NE	Garfield	corn	Ipaga loamy fine sand, 0 to 3 percent slopes	
SRS-2112	MN	Watowan	corn	Darfur fine sandy loam	Coarse-loamy, mixed, superactive, calcareous, mesic Typic Endoaquolls
SRS-2111	CO	Rio Grande	Potato	Norte gravelly sandy loam	Nuetral
SRS-2110	BC	PARC	apples	Typic Haploxeroll	Sediments
SRS-2109	NC	Rowan	Forest / Pasture	Enon fine sandy loam	Fine, mixed, active, thermic Ultic Hapludalfs
SRS-2108	ND	Dickey	corn	Embden fine sandy loam	Coarse-loamy, mixed, superactive, frigid pachic Hapludolls
SRS-2107	OH	Champaign	Corn	Miami silt loam, 2 to 6 percent slopes	Fine-loamy, mixed, active, mesic Oxyaquic Hapludalfs
SRS-2106	WA	Benton	Corn	Shano silt loam	Coarse-silty, mixed, superactive, mesic Xeric Haplocambids
SRS-2105	KY	Breathitt	Corn	Chagrín-Grigsby complex loam, 0 to 6 percent slopes,	Fine-loamy, mixed, active, mesic Dystric Fluventic Eutrudepts
SRS-2104	CA	Madera	Grapes - Rasin	Chino fine sandy loam, 0 to 1	Fine-loamy, mixed, superactive, thermic Aquic Haploxerolls
SRS-2103	ON	Huron	wheat	Loam	
SRS-2102	KS	Kingman	Wheat	Kaski loam, occasionally flooded	Fine-loamy, mixed, superactive, mesic Cumulic Haplustolls
SRS-2101	IL	Iroquois	Pasture	Del Rey silt loam, 0-2 % slopes	Fine, illitic, mesic Aeric Epiaqualls
SRS-2001	WI	Trempleau	pasture	Seaton silt loam, 20 to 30 percent slopes, eroded	Fine-silty, mixed, superactive, mesic Typic Hapludalfs
SRS-2001	CA	Riverside	fallow	Cibola silty clay loam	Fine-silty, mixed, semiactive, mesic Typic Hapludults
SRS-2001	NC	Rowan	Pasture	Lloyd clay loam, 2 to 8 percent slopes, moderately eroded	Fine, kaolinitic, thermic Rhodic Kanhapludults
SRS-2001	AB		Prairie		orthic dark brown chernozems
SRS-2001	ME	Kennebec	Cover Crop	Paxton fine sandy loam	Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts
SRS-2001	KS	Brown	Soybean	Marshall silty clay loam, 5 to 9 percent slopes	Fine-silty, mixed, superactive, mesic Typic Hapludolls
SRS-2001	ON	Middlesex	corn		
SRS-2001	IN	Porter	Soybean	Adrian muck, drained, 0 to 1 percent slopes	Sandy or sandy-skeletal, mixed, euic, mesic Terric Haplosaprists
SRS-2001	OK	Payne	Hay	Pulaski fine sandy loam, 0 to 1 percent slopes, occasionally flooded	Coarse-loamy, mixed superactive ,nnpacid terhmic Udic Ustifluvents
SRS-2001	ID	Power	Wheat	Newdale silt loam 4-12 percent slopes	Coarse-silty, mixed,superactive, frigid Calcicidic Haploxerolls
SRS-2001	IA	Humbolt	Soybean	Nicollet clay	Fine-loamy, mixed, superactive, mesic Aquic Hapludolls
SRS-2001	AB		Wheat		orthic dark brown chernozems
SRS-2001	AL	Limestone	Soybeans	Abernathy silt loam level phase	Fine, kaolinitic, thermic Rhodic Paleudults

SRS-2001	MI	Clinton	Orchard	Marlette loam, 2 to 6 percent slopes	Fine-loamy, mixed, semiactive, mesic Oxyaquic Glossudalfs
SRS-2001	WA	Skagit	potatoes	Mukilteo variant muck	Dysic, mesic Typic Haplohemists
SRS-1915	IA	Badger	Corn	Canisteo clay loam, 0 to 2 percent slopes	Fine-loamy, mixed, superactive, calcareous, mesic Typic Endoaquolls
SRS-1914	QU	Les Chures-de-la-Chaudiere	fallow		
SRS-1913	KS	Harper	wheat	Shellabarger fine sandy loam, 1 to 3 percent slopes	Fine-loamy, mixed, superactive, mesic Udic Argiustolls
SRS-1912	MT	Big Horn	wheat/fallow	Haverson and Lohmiller soils, channeled	Fine-loamy, mixed, superactive, calcareous, mesic Aridic Ustifluvents
SRS-1911	AL	Lee	Wheat	Marvyn loamy sand, 1 to 6 percent slopes	Fine-loamy, kaolinitic, thermic Typic Kanhapludults
SRS-1910	NE	Sarpy	Corn	Monona-Pohocco-Ida silt loams	Fine-silty, mixed, superactive, mesic Typic Eutrudepts
SRS-1909	CT	Hartford	corn	Broadbrook silt loam, 3 to 8 percent slopes	Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts
SRS-1908	AZ	Pinal	Cotton	Casa Grande sandy loam	Fine-loamy, mixed, superactive, hyperthermic Typic Natrargids
SRS-1907	TN	Haywood	Corn	Loring silt loam, 5 to 8 percent slopes, severely eroded	
SRS-1906	SK		Wheat	Mainly Orthic Me sandy loam	Loam
SRS-1905	LA	East Baton Rouge	Grass	DaA—Deerford-Verdun complex, 0 to 2 percent slopes	Fine-silty, mixed, superactive, thermic Albic Glosic Natraqualfs
SRS-1904	IN	Madison	Soybeans	Crosby Silt loam	Fine, mixed, active, mesic Aeric Epiqualfs
SRS-1903	NY	Chautauqua	Hay	Busti silt loam, 3 to 8 percent slopes	Coarse-loamy, mixed, active, nonacid, mesic Aeric Endoaquepts
SRS-1902	ON	Middlesex	Soybean		
SRS-1901	CA	San Luis Obispo	Oats	Cropley clay, 0 to 2 percent slopes, MLRA 14	Fine, smectitic, thermic Aridic Haploxererts
SRS-1815	OR	Marion	Grass Seed	Stayton silt loam, 0 to 7 percent slopes	Medial, mixed, mesic Lithic Haploxerands
SRS-1814	IA	Wright	corn	Brownnton silty clay loam, 0 to 2 percent slopes	Fine, smectitic, calcareous, mesic Vertic Endoaquolls
SRS-1813	SC	Anderson	Pasture	Cecil clay loam, 6 to 10 percent slopes, eroded	Fine, kaolinitic, thermic Typic Kanhapludults
SRS-1812	AB	Division No. 1	Canola		orthic dark brown chernozems
SRS-1811	WV	Raleigh	Pasture	Cookport-Nallen complex, 3 to 8 percent slopes	Fine-loamy, mixed, active, mesic Aquic Fragiudults
SRS-1810	MN	Jackson	Soybean	Nicollet clay loam 1- 3 % slopes	Fine-loamy, mixed, superactive, mesic Aquic Hapludolls
SRS-1809	NM	San Juan	Potato	Doak loam, 1 to 3 percent slopes	Fine-loamy, mixed, active, mesic Typic Haplargids
SRS-1808	IN	Wayne	Corn	Eldean Loam 2 to 6 percent slopes	Fine, mixed, superactive, mesic Typic Hapludalfs
SRS-1807	GA	Jackson	Pasture	Madison sandy loam, 6 to 10 percent slopes	Fine, kaolinitic, thermic Typic Kanhapludults
SRS-1806	ON	Oxford	Soybean		
SRS-1805	NY	Broome	Pasture	Volusia channery silt loam, 8 to 15 percent slopes	Fine-loamy, mixed, active, mesic Aeric Fragiaquepts
SRS-1804	IL	Clay	Soybeans	Bluford silt loam, 0 to 2 percent slopes	Fine, smectitic, mesic Aeric Fragic Epiqualfs
SRS-1803	SC	Anderson	Corn	Cecil sandy loam, 6 to 10 percent slopes	Fine, kaolinitic, thermic Typic Kanhapludults
SRS-1802	NS	Colchester	corn		
SRS-1801	MT	Cascade	wheat	Korent loam , 0 -3 percent	Fine-loamy, mixed, superactive, calcerous frigid mollic Ustifluent
SRS-1715	TX	Nachogdoches	Pasture	Nacogdoches gravely fine sandy loam, 1 to 8 percent slopes	Fine, kaolinitic, thermic Rhodic Paleudalfs
SRS-1714	BC	PARC	apples	Typic Haploxeroll	Stratified Glaciolacustrine
SRS-1713	CA	Sonoma	Grapes	Spreckels loam, 9 to 15 percent slopes	Fine, mixed, superactive, mesic Ultic Palixeralfs
SRS-1712	ME	Aroostok	potato	Linneus silt loam, 0 to 8 percent slopes	Coarse-loamy, isotic, frigid Dystric Eutrudepts
SRS-1711	IA	O'Brien	corn	Sac silty clay loam, 2 to 5 percent slopes	Fine-silty, mixed, superactive, mesic Oxyaquic Hapludolls
SRS-1710	MN	Nobles	soybean	Thurman sandy loam, 2 to 6 percent slopes	Sandy, mixed, mesic Udorthentic Haplustolls
SRS-1709	CA	Sonoma	Grapes	Goulding-Toomes complex, 9 to 50 percent slopes	Loamy, mixed, superactive, thermic Lithic Haploxerepts
SRS-1708	SK	Division #1	Canola	Clay Loam	
SRS-1707	NH	Cheshire	pasture	Colton loamy fine sand, 15 to 50 percent slopes	Sandy-skeletal, isotic, frigid Typic Haplorthods
SRS-1706	GA	Crisp	Hay	Fuquay loamy sand, 0 to 5 percent slopes	Loamy, kaolinitic, thermic Arenic Plinthic Kandiodults
SRS-1705	ID	Fremont	Wheat	Rin silt loams, 1 to 4 percent slopes	Coarse-silty, mixed, superactive Pachic Haplocryolls
SRS-1704	MI	Kalamazoo	corn	Oshtemo sandy loam	Coarse-loamy, mixed, active, mesic Typic Hapludalfs
SRS-1703	TX	Ochiltree	wheat	Darrouzett clay loam, 0 to 1 percent slopes	Fine, mixed, superactive, thermic Pachic Paleustolls
SRS-1702	MA	Franklin	cover crop	Unadilla silt loam, 0 to 3 percent slopes	Coarse-silty, mixed, active, mesic Typic Dystrudepts
SRS-1701	AB	Division #2	Canola		orthic dark brown chernozems
SRS-1615	IN	Terre Haute	Corn	Ade loamy fine sand, 2 to 6 percent slopes	Coarse-loamy, mixed, superactive, mesic Lamellic Argiudolls
SRS-1614	WA	Waitsburg	W wheat	Palouse silt loam,	Fine-silty, mixed, superactive, mesic Pachic Ultic Haploxerolls
SRS-1613	ON	Mount Bridges	Soybean		
SRS-1612	AR	Forest City	-	Calloway silt loam, 1 to 3 percent slopes	Fine-silty, mixed, active, thermic Aquic Fraglossudalfs
SRS-1611	IL	Byron	Corn	Osco silt loam, 2 to 5 percent slopes	Fine-silty, mixed, superactive, mesic Typic Argiudolls
SRS-1610	DE	Sussex	Corn		
SRS-1609	MS	Pike	Grass	RuD3—Ruston fine sandy loam, 8 to 12 percent slopes, severely eroded	Fine-loamy, siliceous, semiactive, thermic Typic Paleudults
SRS-1608	NM	Chaves	Alfalfa	Pecos silty clay loam, 0 to 1 percent slopes	Fine, mixed, superactive, calcareous, thermic Vertic Torrifuvents
SRS-1607	PE	Queens	potato		
SRS-1606	KS	Sherman	corn	Kuma-Keith silt loams, 0 to 2 percent slopes	Fine-silty, mixed, superactive, mesic Pachic Argiustolls
SRS-1605	AL	Lee	corn	Marvyn loamy sand, 1 to 6 percent slopes	Fine-loamy, kaolinitic, thermic Typic Kanhapludults
SRS-1604	CA	Sonoma	grapes	Sebastopol sandy loam, 15-30 percent slopes	Fine, mixed, semiactive, mesic Typic Haploxerults
SRS-1603	ME	Kennebec	fallow	Woodbridge fine sandy loam, 3 to 8 percent slopes	Coarse-loamy, mixed, active, mesic Aquic Dystrudepts
SRS-1602	ON	Huron	clover		
SRS-1601	SD	McCook	soybean	Clarno-Davison loams, 2 to 5 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Haplustolls
SRS-1515	MT	Yellow Stone	wheat/fallow	Shaak silty clay loam, 1 to 4 percent slopes	Fine, smectitic, frigid Vertic Paleustolls
SRS-1514	WI	Iowa	Soybean	Tama silt loam, 0 to 2 percent slopes	Fine-silty, mixed, superactive, mesic Typic Argiudolls

SRS-1513	FL	Alachua	Forest	Millhopper-Urban land complex, 0 to 5 percent slopes	Loamy, siliceous, semiactive, hyperthermic Grossarenic Paleudults
SRS-1512	NE	Buffalo	Corn	Cozad silt loam, 1 to 3 percent slopes	Coarse-silty, mixed, superactive, mesic Typic Haplustolls
SRS-1511	QU	Les Chures-de-la-Chaudiere	buckwheat		
SRS-1510	SC	Anderson	Corn	Pacolet sandy loam, 15 to 25 percent slopes	Fine, kaolinitic, thermic Typic Kanhapludults
SRS-1509	BC	PARC	apples	Typic Haploxeroll	Orthic Dark Brown Chernozem
SRS-1508	AZ	Pinal	Cotton	Trix sandy clay loam	Fine-loamy, mixed, superactive, calcareous, hyperthermic Typic Torrifluvents
SRS-1507	MN	Martin	corn	Blue Earth mucky silty clay loam	Fine-silty, mixed, superactive, calcareous, mesic Mollic Fluvaquents
SRS-1506	CT	Tolland	pasture	Agawam fine sandy loam, 3 to 8 percent slopes	Coarse-loamy over sandy or sandy-skeletal, mixed, active, mesic Typic Dystrudepts
SRS-1505	IA	Hardin	corn	Webster-Nicollet complex, 1 to 3 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Endoaquolls
SRS-1504	ID	Jerome	Potato	Purdam silt loam, 1 to 4 percent slopes	Fine-silty, mixed, superactive, mesic Haploxeralfic Argidurids
SRS-1503	TX	Nachogdoches	Pasture	Darco loamy fine sand, 1 to 8 percent slopes	Loamy, siliceous, semiactive, thermic Grossarenic Paleudults
SRS-1502	KS	Brown	Soybean	Wymore silty clay loam, 3 to 6 percent slopes	Fine, smectitic, mesic Aquertic Argiudolls
SRS-1501	NS	Colchester	corn		
SRS-1415	MT	Judith Basin	wheat	Danvers-Judith clay loams, 0 to 2 percent slopes	Fine-loamy, carbonatic, frigid Typic Calcicustolls
SRS-1414	NE	Cass	Soybeans	Otoe silty clay loam, 6 to 11 percent slopes, eroded	Fine, smectitic, mesic Aquertic Hapludalfs
SRS-1413	IL	Iroquois	timber	Del Rey silt loam 0-2 % slope	Fine, illitic, mesic Aerlic Epiaqualls
SRS-1412	OK	Payne	Hay	Pulaski fine sandy loam, 0 to 1 percent slopes, occasionally flooded	Coarse-loamy, mixed, superactive, nonacid, thermic Udic Ustifluvents
SRS-1411	VT	Windham	pasture	Warwick channery fine sandy loam	Loamy-skeletal, mixed, active, mesic Typic Dystrudepts
SRS-1410	IN	Randolph	Soybean	Saranac silty clay, frequently flooded	Fine, mixed, active, mesic Fluvaquentic Endoaquolls
SRS-1409	IA	Palo Alto	corn	Canisteo silty clay loam, 0 to 2 percent slopes	Fine-loamy, mixed, superactive, calcareous, mesic Typic Endoaquolls
SRS-1408	MD	Prince Georges	soybean	Elkton silt loam, 0 to 2 percent slopes	Fine-silty, mixed, active, mesic Typic Endoaquolls
SRS-1407	GA	Grady	Onions	Tifton loamy sand, 2 to 5 percent slopes	Fine-loamy, kaolinitic, thermic Plinthic Kandiodults
SRS-1406	AZ	Mariacopa	Forage Grass	Brios sandy loam	Sandy, mixed, hyperthermic Typic Torrifluvents
SRS-1405	CA	Fresno	Native	Delhi loamy sand, 0 to 3 percent slopes	Mixed, thermic Typic Xeropsamments
SRS-1404	TN	Wilson	Pasture	Bradyville silt loam, 2 to 5 percent slopes, eroded	Fine, mixed, semiactive, thermic Typic Hapludalfs
SRS-1403	SD	Brookings	Soybeans	Lanona-Swenoda sandy	Coarse-loamy, mixed, superactive, frigid Calcic Hapludolls
SRS-1402	NS	Westmoreland	Pasture		?
SRS-1401	MO	Boone	Alfalfa	Arisburg silt loam, 1 to 3 percent slopes	Fine, smectitic, mesic Aquertic Argiudolls
SRS-1315	AR	Marianna	Misc	Calloway silt loam, 0 to 1 percent slopes	Fine-silty, mixed, active, thermic Aquic Fraglossudalfs
SRS-1314	WY	Sheridan	Pasture		
SRS-1313	WI	Woods	Pasture		
SRS-1312	ME	Orno	small grains	Howland very stony loam, 0 to 8 percent slopes	Coarse-loamy, isotic, frigid Aquic Haplorthods
SRS-1311	MT	Creston	barley	Creston silt loam, 0 to 3 percent slopes	Fine-silty, mixed, superactive Typic Haploborolls
SRS-1310	SK	Division #1	Wheat	Cherazom	
SRS-1309	NM	San Juan	Corn	Shiprock fine sandy loam, 2 to 5 percent slopes	Coarse-loamy, mixed, superactive, mesic Typic Haplargids
SRS-1308	AL	Limestone	Soybeans	Decatur silty clay loam eroded rolling phase	Fine, kaolinitic, thermic Rhodic Paleudults
SRS-1307	NJ	Salem	Veg	Mattapex silt loam, 2 to 5 percent slopes	Fine-silty, mixed, active, mesic Aquic Hapludults
SRS-1306	IN	Pulaski	Corn	TmaAN—Toto muck, drained, 0 to 1 percent slopes	Coprogenous, euic, mesic Limnic Haplosaprists
SRS-1305	WA	Benton	Wheat	Ritzville silt loam	Coarse-silty, mixed, superactive, mesic Calcicidic Haploxerolls
SRS-1304	IA	Boone	soybeans	Canisteo silty clay loam, 0 to 2 percent slopes	Fine-loamy, mixed, superactive, calcareous, mesic Typic Endoaquolls
SRS-1303	RI	Oak Island	Peaches	Paxton very stony fine sandy loam	Coarse-loamy, mixed, active, mesic Oxyaquic Dystrudepts
SRS-1302	MI	Branch	Soybean	Elmdale fine sandy loam, 2 to 6 percent slopes	Coarse-loamy, mixed, semiactive, mesic Oxyaquic Hapludalfs
SRS-1301	FL	Alachua	Forest	Tavares sand, 0 to 5 percent slopes	Hyperthermic, uncoated Typic Quartzipsamments
SRS-1215	MT	Big Timber	Oats	Farnuf Loam	Fine-loamy, mixed, superactive, frigid Typic Argiustolls
SRS-1214	NE	Sarpy	Corn	Contrary-Monona-Ida complex, 6 to 17 percent slopes	Fine-silty, mixed, superactive, mesic Typic Hapludolls
SRS-1213	CA	Fresno	Cotton	Fresno Sandy Loam	
SRS-1212	MS	Yahzoo	Corn	Dundee Silt Loam 0-2% slopes	Fine-silty, mixed, active, thermic Typic Endoaqualls
SRS-1211	PA	Erie	Grass	Mardin silt loam, 3 to 8 percent slopes	Coarse-loamy, mixed, active, mesic Typic Fragiudepts
SRS-1210	WA	Benton	Potato	Quincy loamy fine sand	Mixed, mesic Xeric Torripsamments
SRS-1209	IA	Boone	Pasture	Hayden loam, 2 to 5 percent slopes	Fine-loamy, mixed, superactive, mesic Glossic Hapludalfs
SRS-1208	LA	Acadia	Rice	Crowley silt loam, 0 to 1 percent slopes	Fine, smectitic, thermic Typic Albaqualls
SRS-1207	NH	Rockingham	Pasture	Deerfield fine sandy loam, 3 to 8 percent slopes	Mixed, mesic Aquic Udipsamments
SRS-1206	IN	Jasper	Soybeans	Chelsea sand, 2 to 6 percent slopes	Mixed, mesic Lamellic Udipsamments
SRS-1205	MB	Cromer	small grains		
SRS-1204	VA	Chesterfield	small grains	Appling- Spotsylvania sandy loams, 2 to 6 percent slopes	Fine, kaolinitic, thermic Typic Kanhapludults
SRS-1203	IA	Story City	soybeans	Nicollet loam, 1 to 3 percent slopes	Fine-loamy, mixed, superactive, mesic Aquic Hapludolls
SRS-1202	NY	Tully	hay	Palmyra gravelly loam	Sandy loam
SRS-1201	NV	Winnemeca	potatoes	Batan-Goldrun-Bubus complex, 0 to 30 percent slopes	Fine-silty, mixed, superactive, calcareous, mesic Duric Torriorthents
SRS-1115	UT	Plymouth	wheat	Hansel silt loam, 1 to 6 percent slopes	Fine-silty, mixed, superactive, mesic Calcic Haploxeralfs
SRS-1114	IL	McClean	corn	Chenoa silty clay loam, 2 to 5 percent slopes	Fine, illitic, mesic Aquic Argiudolls
SRS-1113	NE	Kearney	corn	Holdrege-Hall silt loams, 0 to 1 percent slopes	Fine-silty, mixed, superactive, mesic Typic Argiustoll
SRS-1112	MA	Franklin	potatoes	Winooski silt loam, 0 to 3 percent slopes, protected	Coarse-silty, mixed, superactive, mesic Fluvaquentic Dystrudepts
SRS-1111	AL	Lee	cotton	Marvyn loamy sand	Fine-loamy, kaolinitic, thermic Typic Kanhapludults
SRS-1110	ND	Grand Forks	wheat	Lowe loam, channeled, 0 to 2 percent slopes	Fine-loamy, mixed, superactive, frigid Typic Calciaqualls

SRS-1109	ID	Cassia	wheat	Davey fine sandy loam, 0 to 2 percent slopes	Sandy, mixed, mesic Xeric Haplocambids
SRS-1108	NC	Currituck	corn	Roanoke fine sandy loam	Fine, mixed, semiactive, thermic Typic Endoaquults
SRS-1107	CT	Storrs	pasture	Woodbridge fine sandy loam, 0 to 3 percent slopes	Coarse-loamy, mixed, active, mesic Aquic Dystrudepts
SRS-1106	WA	Plymouth	potatoes	Warden silt loam, 0 to 5 percent slopes	Coarse-silty, mixed, superactive, mesic Xeric Haplocambids
SRS-1105	IA	Webster	corn	Webster silty clay loam, 0 to 2 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Endoaquolls
SRS-1104	CA	Dehli	almonds	Atwater loamy sand, deep over hardpan, 0 to 3 percent slopes	Coarse-loamy, mixed, active, thermic Typic Haploxerafals
SRS-1103	TN	Lebanon	corn	Bradyville silt loam, 2 to 5 percent slopes, eroded	Fine, mixed, semiactive, thermic Typic Hapludalfs
SRS-1102	MI	Cass	soybeans	Kalamazoo loam, 0 to 2 percent slopes	Fine-loamy, mixed, semiactive, mesic Typic Hapludalfs
SRS-1101	PEI	Harrington, PEI	small grains	fine sandy loam soil	Orthic Humo-Ferric Podzol
SRS-1015	IA	Britt	corn	Clarion loam, 2 to 5 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Hapludolls
SRS-1014	CA	San Joaquin	cotton	Tachi clay, 0 to 1 percent slopes	Very-fine, smectitic, thermic Typic Natraquerts
SRS-1013	FL	Callahan	pasture	Surrency loamy fine sand	Loamy, siliceous, semiactive, thermic Arenic Umbric Paleaquults
SRS-1012	WV	Petersburg	hay	Tioga Fine sandy loam	Coarse-loamy, mixed, superactive, mesic Dystric Fluventic Eutrudepts
SRS-1011	IL	Cropsey	corn	Chenoa silty clay loam, 2 to 5 percent slopes	Fine, illitic, mesic Aquic Argiudolls
SRS-1010	WI	Trempealeu	corn	Gotham Loamy Fine Sand	Mixed, mesic Psammentic Hapludalfs
SRS-1009	TX	Bell	corn	Frio Silty Clay 0-1% Slope	Fine, smectitic, thermic Cumulic Haplustolls
SRS-1008	AR	Washington	pasture	Captina silt loam, 1 to 3 percent slopes	Fine-silty, siliceous, active, mesic Typic Fragiudults
SRS-1007	IA	Kossuth	soybeans	Webster Silty Clay Loam 0-2% slopes	Fine-loamy, mixed, superactive, mesic Typic Endoaquolls
SRS-1006	SC	Anderson	corn	Cartecay-Chewacla Complex	Coarse-loamy, mixed, semiactive, nonacid, thermic Aquic Udifluvents
SRS-1005	SD	Brookings	corn	Lanona-Swenoda sandy loams, 2 to 6 percent slopes	Coarse-loamy, mixed, superactive, frigid Calcic Hapludolls
SRS-1004	IL	Livingston	corn	Chenoa silty clay loam, 0 to 2 percent slopes	Fine, illitic, mesic Aquic Argiudolls
SRS-1003	CA	Sonoma	grapes	Goldridge fine sandy loam--on a south facing slope of 5 percent	Fine-loamy, mixed, superactive, isomesic Typic Haplustults
SRS-1002	ID	Jerome	pasture	Chiara silt loam, 1 to 8 percent slopes	Loamy, mixed, superactive, mesic, shallow Xeric Haplodurids
SRS-1001	KY	Daviess	soybean	Weinbach silt loam	Fine-silty, mixed, active, mesic Aeric Fragiaqualfs
SRS-0915	AL	Lee	cotton	Marvyn loamy sand	Fine-loamy, kaolinitic, thermic Typic Kanhapludults
SRS-0914	NE	Holt	corn	Valentine- Dunday loamy fine sands, 3 to 9 percent slopes	Mixed, mesic Typic Ustipsammets
SRS-0913	IA	Hancock	corn	Nicolette clay loam	Fine-loamy, mixed, superactive, mesic Aquic Hapludolls
SRS-0912	CA	Madera	cotton	Chino loam, slightly saline-alkali, 0 to 1 percent slopes	Fine-loamy, mixed, superactive, thermic Aquic Haploxerolls
SRS-0911	MN	Stearns	soybeans	Waukon loam, 2 to 6 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Argiaquolls
SRS-0910	ID	Power	potatoes	Penoyer silt loam, 0 to 2 percent slopes	Coarse-silty, mixed, superactive, calcareous, mesic Typic Torriorthents
SRS-0909	IA	Wright	soybeans	Harps clay loam, 0 to 2 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Calciaquolls
SRS-0908	KS	Harper	wheat	Nalim loam, 0 to 1 percent slopes	Fine-loamy, mixed, superactive, mesic Udic Argiustolls
SRS-0907	IL	Kankakee	corn	Darroch silt loam, 0 to 2 percent slopes	Fine-loamy, mixed, mesic Aquic Argiudolls
SRS-0906	GA	Crisp	cotton	Orangeburg loamy sand, 2 to 5 percent slopes	Fine-loamy, kaolinitic, thermic Typic Kandiuults
SRS-0905	IN	Fountain	soybeans	Mahalalad silty clay loam, 0 to 1 percent slopes	Fine-silty, mixed, superactive, mesic Typic Argiaquolls
SRS-0904	VA	Chesterfield	pasture	Bourne fine sandy loam, 2 to 6 percent slopes	Fine-loamy, mixed, semiactive, thermic Typic Fragiudults
SRS-0903	IA	Wright	corn	Ottosen clay loam, 1 to 3 percent slopes	Fine-loamy, mixed, superactive, mesic Aquic Hapludolls
SRS-0902	NM	Dona Ana	pecans	Glendale clay loam	Fine-silty, mixed, superactive, calcareous, thermic Typic Torrifluvents
SRS-0901	ID	Fremont	pasture	Marystown silt loam, 1 to 4 percent slopes	Fine-silty, mixed, superactive, frigid Pachic Argixerolls
SRS-0815	IL	Kankakee	corn	Reddick clay loam, 0 to 2 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Endoaquolls
SRS-0814	CO	Logan	corn	Loveland Clay Loam	Fine-loamy over sandy or sandy-skeletal, mixed, superactive, calcareous, mesic Fluvaquentic
SRS-0813	AZ	Pinal	vegetables	Casa Grande sandy loam, 0-3 percent slope	Fine-loamy, mixed, superactive, hyperthermic Typic Natrargids
SRS-0812	IA	Buchanan	soybeans	Schley loam, 1 to 4 percent slopes	Fine-loamy, mixed, superactive, mesic Udollic Endoaqualfs
SRS-0811	NE	Sarpy	soybeans	Contrary- Monona silty clay loams, 6 to 11 percent slopes	Fine-silty, mixed, superactive, mesic Typic Hapludolls
SRS-0810	NE	Holt	corn	Dunday loamy sand, 0 to 3 percent slopes	Sandy, mixed, mesic Entic Haplustolls
SRS-0809	IA	Hamilton	corn	Clarion loam, 2 to 5 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Hapludolls
SRS-0808	ID	Bingham	potatoes	Bock loam, 0 to 2 percent slopes	Coarse-loamy, mixed, superactive, frigid Calcic Haploxerolls
SRS-0807	OR	Polk	small grains	Woodburn silt loam, 0 to 3 percent slopes	Fine-silty, mixed, superactive, mesic Aquultic Argixerolls
SRS-0806	WI	Portage	corn	Rozellville loam 0-3 percent slopes	Fine-loamy, mixed, superactive, frigid Haplic Glossudalfs
SRS-0805	MN	Swift	soybeans	Bearden- Quam, depressional, complex, 0 to 2 percent slopes	Fine-silty, mixed, superactive, frigid Aeric Calciaquolls
SRS-0804	IL	Ford	soybeans	Elliott silty clay loam, 2 to 4 percent slopes, eroded	Fine, illitic, mesic Aquic Argiudolls
SRS-0803	MN	Brown	corn	Linder sandy loam	Coarse-loamy, mixed, superactive, mesic Aquic Hapludolls
SRS-0802	OK	Payne	small grains	Teller fine sandy loam 0- 3 percent slopes	Fine-loamy, mixed, active, thermic Udic Argiustolls
SRS-0801	NE	Holt	corn	Dunday loamy sand, 0 to 3 percent slopes	Sandy, mixed, mesic Entic Haplustolls
SRS-0715	IA	Cerro Gordo	soybeans	Clarion loam, 2 to 5 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Hapludolls
SRS-0714	CA	Tulare	cotton	Colpien loam, 0 to 2 percent slopes	Fine-loamy, mixed, superactive, thermic Calcic Pachic Haploxerolls
SRS-0713	WA	Adams	potatoes	Shano silt loam, 0 to 5 percent slopes	Coarse-silty, mixed, superactive, mesic Xeric Haplocambids
SRS-0712	IN	Hendricks	corn	Whitaker silt loam, 0-3 percent slopes	Fine-loamy, mixed, active, mesic Aeric Endoaqualfs
SRS-0711	OH	Preble	corn	Miamian- Losantville clay loams, 6 to 12 percent slopes, severely eroded	Fine, mixed, active, mesic Oxyaquic Hapludalfs
SRS-0710	MN	Houston	corn	Seaton silt loam, valleys, 6 to 12 percent slopes, eroded	Fine-silty, mixed, superactive, mesic Typic Hapludalfs
SRS-0709	MN	Swift	soybeans	Bearden- Quam, depressional, complex, 0 to 2 percent slopes	Fine-silty, mixed, superactive, frigid Aeric Calciaquolls
SRS-0708	NE	Saline	corn	Crete silt loam, 1 to 3 percent slopes	Fine, smectitic, mesic Pachic Argiustolls
SRS-0707	CO	Rio Blanco	forest	Routt loam, 3 to 25 percent slopes	Fine, smectitic Pachic Palecryolls
SRS-0706	WI	Jefferson	soybeans	Grellton fine sandy loam, 2 to 6 percent slopes	Fine-loamy, mixed, superactive, mesic Typic Hapludalfs

SRS-0705	CA	Napa	grapes	Pleasanton gravelly fine sandy loam	Fine-loamy, mixed, superactive, thermic Mollic Haploxerafals
SRS-0704	WA	Grant	corn	Ephrata fine sandy loam, 2 to 5 percent slopes	Coarse-loamy over sandy or sandy-skeletal, mixed, superactive, mesic Xeric Haplocambids
SRS-0703	NE	Holt	corn	Dunday loamy sand, 0 to 3 percent slopes	Sandy, mixed, mesic Entic Haplustolls
SRS-0702	IA	Iowa	soybeans	Otley silty clay loam, 2 to 5 percent slopes	Fine, smectitic, mesic Oxyaquic Argiudolls
SRS-0701	WI	Trempaleau	corn	Gotham Loamy Fine Sand	Mixed, mesic Psammentic Hapludalfs
SRS-0605	WA	Grant	wheat	Ephrata fine sandy loam, 2 to 5 percent slopes	Coarse-loamy over sandy or sandy-skeletal, mixed, superactive, mesic Xeric Haplocambids
SRS-0604	TX	Bell	corn	Frio Silty Clay 0-1% Slope	Fine, smectitic, thermic Cumulic Haplustolls
SRS-0603	MN	Houston	corn	Seaton silt loam, valleys, 6 to 12 percent slopes, eroded	Fine-silty, mixed, superactive, mesic Typic Hapludalfs
SRS-0602	IL	Kankakee	soybeans	Andres silt loam, 0 to 2 percent slopes	Fine-loamy, mixed, superactive, mesic Aquic Argiudolls
SRS-0601	NE	Holt	corn	Dunday loamy sand, 0 to 3 percent slopes	Sandy, mixed, mesic Entic Haplustolls